

DIRECTIONS FOR REMOVAL:

Within 24 hours, soak GlykoSL[®] with water and gently remove.

Store at controlled room temperature,
15 °C to 30 °C (59 °F to 86 °F).

Sterilized with radiation.

Do not resterilize or reuse.

Single use only.

Not Made with Natural Rubber Latex



Emergo Europe
Molenstraat 15
2513 BH, The Hague
The Netherlands



Marine Polymer
Technologies, Inc.
107 Water Street
Danvers, MA 01923
1-888-666-2560

GlykoSL[®]

REF 400-13

Distributed by:
Marine Polymer Technologies, Inc.
107 Water Street • Danvers, MA 01923
1-888-666-2560
Made in the U.S.A.

MPS 500-28 rev 04

CAUTION: Federal (USA) law restricts this device to sale by or on the order of a physician or other licensed practitioner.

DESCRIPTION:

GlykoSL® is a conformable, soft and absorbent calcium alginate dressing attached to a foam backing.

INDICATIONS:

GlykoSL® is indicated for use in the local management of bleeding wounds such as vascular access sites and percutaneous catheters or tubes.

PRECAUTIONS:

There are no known contraindications to the use of GlykoSL®. Longer compression may be necessary for hypertensive or obese patients.

DIRECTIONS FOR APPLICATION:

1. Open GlykoSL® pouch into sterile field.
2. Hold pressure proximal to the puncture site.
3. Place GlykoSL® over puncture site, **smooth side away from skin.**
4. Apply firm compression over the puncture site and release the proximal pressure after 3 to 4 minutes.
5. Hold pressure on puncture site until hemostasis is achieved, then slowly release.
6. Place dry gauze over GlykoSL® and cover with a transparent dressing.