

**Important: Read all product information before using. Keep the box for important information. This product is intended for use in children.**

**Active ingredient (in each 5 mL = 1 teaspoonful)**

Ibuprofen 100 mg (NSAID)\*.....Pain reliever/fever reducer

\*nonsteroidal anti-inflammatory drug

**Uses** temporarily: ■ relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache ■ reduces fever

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if your child: ■ has had stomach ulcers or bleeding problems ■ takes a blood thinning (anticoagulant) or steroid drug ■ takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ takes more or for a longer time than directed. **Sore throat warning:** Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor. **Do not use**

■ if the child has ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery. **Ask a doctor before use if**

■ stomach bleeding warning applies to your child ■ child has a history of stomach problems, such as heartburn ■ child has problems or serious side effects from taking pain relievers or fever reducers ■ child has not been drinking fluids ■ child has lost a lot of fluid due to vomiting or diarrhea ■ child has high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ child has asthma ■ child is taking a diuretic. **Ask a doctor or pharmacist before use if the child is** ■ under a doctor's care for any serious condition ■ taking any other drug. **When using this product**

■ give with food or milk if stomach upset occurs ■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed. **Stop use and ask a doctor if** ■ child experiences any of the following signs of stomach bleeding ■ feels faint ■ vomits blood ■ has bloody or black stools ■ has stomach pain that does not get better ■ the child does not get any relief within first day (24 hours) of treatment ■ fever or pain gets worse or lasts more than 3 days

■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child: ■ has had stomach ulcers or bleeding problems ■ takes a blood thinning (anticoagulant) or steroid drug ■ takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ takes more or for a longer time than directed. **Sore throat warning:** Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor. **Do not use** ■ if the child has ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery. **Ask a doctor before use if** ■ stomach bleeding warning applies to your child ■ child has a history of stomach problems, such as heartburn ■ child has problems or serious side effects from taking pain relievers or fever reducers ■ child has not been drinking fluids ■ child has lost a lot of fluid due to vomiting or diarrhea ■ child has high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ child has asthma ■ child is taking a diuretic. **Ask a doctor or pharmacist before use if the child is** ■ under a doctor's care for any serious condition ■ taking any other drug. **When using this product** ■ give with food or milk if stomach upset occurs ■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed. **Stop use and ask a doctor if** ■ child experiences any of the following signs of stomach bleeding ■ feels faint ■ vomits blood ■ has bloody or black stools ■ has stomach pain that does not get better ■ the child does not get any relief within first day (24 hours) of treatment ■ fever or pain gets worse or lasts more than 3 days

NDC 0904-5577-20

 **MAJOR**<sup>®</sup>  
Children's  
**Ibuprofen**

**Oral Suspension**

100 mg per 5 mL

Pain Reliever/Fever Reducer (NSAID)

**Lasts up to 8 hours**

**For Ages  
2 to 11  
Years**



**Grape Flavored Liquid**

Alcohol Free

**120 mL (4 FL OZ)**

■ redness or swelling is present in the painful area ■ any new symptoms appear. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)  
**Directions** ■ this product does not contain directions or complete warnings for adult use ■ do not give more than directed ■ shake well before using ■ mL = milliliter; tsp = teaspoonful ■ find right dose on chart. If possible, use weight to dose; otherwise use age. ■ use only enclosed dosing cup. Do not use any other dosing device. ■ if needed, repeat dose every 6-8 hours ■ do not use more than 4 times a day ■ replace original bottle cap to maintain child resistance ■ wash dosage cup after each use

**Dosing Chart**

Weight (lb)	Age (yr)	Dose (mL or tsp)**
under 24 lbs	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL (1 tsp)
36-47 lbs	4-5 years	7.5 mL (1½ tsp)
48-59 lbs	6-8 years	10 mL (2 tsp)
60-71 lbs	9-10 years	12.5 mL (2½ tsp)
72-95 lbs	11 years	15 mL (3 tsp)

\*\*or as directed by a doctor

**Other information** ■ each 5 mL (1 tsp) contains: sodium 2 mg

■ do not use if printed neckband is broken or missing

■ store at 20-25°C (68-77°F) ■ do not freeze

**Inactive ingredients** anhydrous citric acid, D&C red #33, FD&C blue #1, FD&C red #40, glycerin, high fructose corn syrup, hypromellose, natural and artificial grape flavor, polysorbate 80, purified water, sodium benzoate, sorbitol solution, xanthan gum

**Questions or comments?** 1-800-616-2471

Distributed by  
**MAJOR**<sup>®</sup> PHARMACEUTICALS

31778 Enterprise Drive

Livonia, MI 48150 USA

M-05 REV. 11/15

Re-Order No. 005979

**CODE  
AREA**

**Gluten Free** • no varnish • no color  
: 66026 5C F8