

Drug Facts

Active ingredient (in each tablet)

Naproxen sodium 220 mg
(naproxen 200 mg (NSAID))fever reducer
nonsteroidal anti-inflammatory drug

Purposes

■ temporarily relieves minor aches and pains due to:
■ minor pain of arthritis ■ muscular aches ■ backache
■ menstrual cramps ■ headache ■ toothache
■ the common cold
■ temporarily reduces fever


Warnings

Always alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
■ hives ■ facial swelling ■ rash ■ wheezing
■ stomach pain ■ skin redness ■ asthma ■ blisters
If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older
■ have had stomach ulcers or bleeding problems
■ take a blood thinning anticoagulant or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product
■ take more or for a longer time than directed

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery

Ask a doctor before use if ■ the stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have asthma
Ask a doctor a pharmacist before use if you are
■ under a doctor's care in any serious condition
■ taking any other drug



CODE AREA
LOT NO. 5
EXP. 053611094063

Drug Facts (continued)

When using this product
■ take with food or milk if stomach upset occurs
■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding:
■ feel faint ■ vomit blood ■ have bloody or black stools
■ have stomach pain that does not get better
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ you have difficulty swallowing
■ it feels like the pill is stuck in your throat
■ redness or swelling is present in the painful area
■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions


■ do not take more than directed
■ the smallest effective dose should be used
■ drink a full glass of water with each dose
Adults and children 12 years and older:
■ take 1 tablet every 8 to 12 hours while symptoms last
■ for the first dose you may take 2 tablets within the first hour
■ do not exceed 2 tablets in any 8- to 12-hour period
■ do not exceed 3 tablets in a 24-hour period
Children under 12 years: ■ ask a doctor

Other information

■ each tablet contains: sodium 21 mg
■ store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Gluten Free
***This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Aleve®.
Distributed by: Rugby Laboratories
3778 Enterprise Drive
Livonia, MI 48150
www.rugbylaboratories.com
Re-Order No. 370445 REV. 06/15 R-05
Questions or comments? 1-800-645-2158
Inactive ingredients FD&C blue no. 2 aluminum powder, talc, titanium dioxide
Inactive ingredients FD&C blue no. 2 aluminum microcrystalline cellulose, polyethylene glycol, lake, hypromellose, magnesium stearate, croscarmellose, polyethylene glycol.

DO NOT USE IF PRINTED FOIL UNDER CAP IS BROKEN OR MISSING



ACTUAL SIZE



COMPARE to the active ingredient in ALEVE®**

NDC 0536-1094-06

Rugby®

All Day Relief

**Naproxen Sodium Tablets, USP 220 mg
Pain Reliever/Fever Reducer (NSAID)**

Strength to Last 12 Hours

50 TABLETS

ACTUAL SIZE

KEEP CARTON FOR REFERENCE

EFB 110
(490) 490-008 EFB110-2