



NDC 0536-1053-05

ASPIRIN

Pain Reliever/
Fever Reducer
(NSAID)

325 mg each

500 Tablets



AID FOR
RELIEF OF
SIMPLE HEADACHE,
COLD DISCOMFORTS,
MUSCULAR ACHES

Drug Facts

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)*Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Purpose

Uses temporarily relieves: ■ pain and fever of colds ■ headache
■ toothache ■ muscle pain ■ minor pain of arthritis ■ menstrual pain

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older
■ take a blood thinning (anticoagulant) or steroid drug
■ have 3 or more alcoholic drinks every day while using this product
■ take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others] ■ take more or for a longer time than directed
■ have had stomach ulcers or bleeding problems

Do not use

■ if you are allergic to aspirin or any other pain reliever/fever reducer
■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if ■ you have asthma ■ you are taking a diuretic

■ stomach bleeding warning applies to you
■ you have a history of stomach problems, such as heartburn
■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ➔

No print/No varnish
Lot & Exp date

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking a prescription drug for
■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools
■ have stomach pain that does not get better
■ an allergic reaction occurs. Seek medical help right away.
■ ringing in the ears or loss of hearing occurs ■ new symptoms occur
■ pain gets worse or lasts more than 10 days ■ redness or swelling is present
■ fever gets worse or lasts more than 3 days

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

■ drink a full glass of water with each dose
■ adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours
■ children under 12 years: ask a doctor

Other information

■ use by expiration date on package
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Inactive ingredient corn starch

Questions or comments?

1-800-645-2158

50844

REV0915A24914



3 05361 05305 5

Distributed by: Rugby Laboratories, 31778 Enterprise Drive,
Livonia, MI 48150 www.rugbylaboratories.com
R-17 Re-order No. 370396 Rev. 02/16

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**