

## Drug Facts (continued)

### Directions

- do not take more than 6 doses in any 24 hour period
  - use only with enclosed dosing cup. Do not use with any other device.
- |                                       |  |
|---------------------------------------|--|
| adults and children 12 years and over | 10 - 20mL every 4 hours or as directed by doctor |
| children under 12 years               | do not use                                       |

### Other information

- each 5 mL contains: **sodium 3 mg**
- store at 20°-25°C (68°-77°F). Do not refrigerate


**Inactive ingredients** caramel powder, citric acid, corn syrup, FD&C Red #40, glycerin, menthol, propylene glycol, purified water, saccharin sodium, sodium benzoate

**Questions or comments?** (800) 616-2471

Distributed by:

**MAJOR PHARMACEUTICALS**  
31778 Enterprise Drive  
Livonia, MI 48150, USA

M-64 REV 02/16  
Re-Order No. 263921



\* This product is not manufactured or distributed by Wyeth Consumer Healthcare, distributor of Robitussin®. Robitussin® is a registered trademark of A. H. Robins Company, Inc.

**TAMPER-EVIDENT:** Do not use this product if inner foil seal over the mouth of the bottle is cut, torn, broken or missing.

NDC 0904-0061-00

 **MAJOR®**

**ROBAFEN™**  
**cough formula**  
Expectorant

Compare to active ingredient of  
Robitussin® Mucus+Chest Congestion\*.

Guaifenesin USP 100 mg per 5 mL  
**Makes Coughs More Productive**

**4 FL. OZ. (118 mL)**

## Drug Facts

**Active Ingredient (in each 5 mL)** ..... **Purpose**  
Guaifenesin USP 100 mg ..... Expectorant

### Uses

Helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive

### Warnings

**Ask a doctor before use if you have**

- a cough that lasts or a chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- a cough occurs with too much phlegm (mucus)

**Stop use and ask a doctor** if cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use. **Keep out of the reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

PEEL HERE