

Supplement Facts	
Serving Size: 1 teaspoonful (5 mL)	
Servings per container: 94	
Amount per teaspoonful (5 mL)	% Daily Value*
Ferrous sulfate 220 mg (equivalent to 44 mg iron)	240%
* % Daily Value is based on 2000 calorie diet	

Other ingredients: purified water, sucrose, alcohol (5%), sorbitol solution, propylene glycol, citric acid, lemon flavor, sodium benzoate, ascorbic acid, FD&C yellow no. 6, saccharin sodium.

Uses: Iron supplements are used to treat or prevent low blood levels of iron, such as for anemia or during pregnancy.

Warnings

Iron containing products should only be taken for very specific reasons and only under the advice and supervision of a doctor. Consult your doctor for guidance for iron supplementation.
Do not exceed recommended dosage.

Warning:
Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children younger than 6 years of age. Keep this product out of the reach of children. In case of accidental overdose, call a doctor or Poison Control Center immediately.

Do NOT use if • you are allergic to any ingredient in Ferrous Sulfate Elixir • you have high levels of iron in your blood



N 3 0904-1465-16 9

MAJOR®

NDC 0904-1465-16

FeroSul™
Ferrous Sulfate Elixir

Contains Alcohol 5%

Ferrous Sulfate
220 mg/5 mL
(equivalent to 44 mg iron)

**BULK CONTAINER - NOT
FOR HOUSEHOLD USE**

This container is not
child-resistant.
Pharmacist - Dispense in a
child-resistant container.

**DO NOT ACCEPT IF IMPRINTED
SAFETY SEAL AROUND CAP IS
BROKEN OR MISSING**

473 mL (16 fl oz)

Ask a doctor before use if • you are pregnant, planning to become pregnant, or are breast-feeding • you are taking any prescription or over-the-counter medicines

Since oral iron products interfere with absorption of oral tetracycline antibiotics, these products should not be taken within two hours of each other. Occasional gastrointestinal discomfort (such as nausea) may be minimized by taking with meals. Iron containing medications may occasionally cause constipation or diarrhea, and may cause temporary staining of the teeth. Mixing Ferrous Sulfate Elixir with water or fruit juice may help to decrease the effects.

Do not take Ferrous Sulfate Elixir within 1 hour before or 2 hours after antacids, eggs, milk, milk byproducts, coffee, or tea.

Directions

Do not exceed recommended dosage. Do not mix with milk, milk byproducts, or wine-based vehicles.

Adults	1 teaspoonful daily or as directed by a doctor
Children under 12 years	Consult a doctor

Other Information

Store at room temperature 20°-25°C (68°-77°F). Protect from Freezing.

To report adverse drug events call: 800-616-2471

Re-order No. 245837 Rev. 07/09 M-52

Distributed By
MAJOR PHARMACEUTICALS
31778 Enterprise Drive,
LIVONIA, MI 48150, USA