

Adhesive Area

STOP PEELING

Questions or comments?
Call: (800) 616-2471

Drug Facts (continued)

Stop use and ask a doctor if

- you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

■ take with a glass of water	
adults and children 12 years and over	take 1 to 3 tablets in a single daily dose
children 6 to under 12 years of age	take 1 tablet in a single daily dose
children under 6 years of age	ask a doctor


Other information

- each tablet contains: magnesium 5 mg
- store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package
- protect from excessive humidity

Inactive ingredients

acacia, anhydrous calcium sulfate, anhydrous lactose, carnauba wax, colloidal silicon dioxide, corn starch, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, gelatin, iron oxide, iron oxide black, iron oxide yellow (iron oxide ochre), magnesium stearate, microcrystalline cellulose, polyethylene glycol (PEG) 400, polyvinyl acetate phthalate, povidone, stearic acid, sodium starch glycolate, stearic acid, sugar, talc, titanium dioxide

NDC 0904-7927-60



MAJOR®
BISACODYL
Stimulant Laxative

For Temporary Relief of Occasional Constipation

ENTERIC COATED

BISACODYL TABLETS,
USP 5 mg each

100 Tablets

Compare to the active ingredient of DULCOLAX®*

*This product is not manufactured or distributed by Boehringer Ingelheim Pharmaceuticals, Inc., owner of the registered trademark Dulcolax®. 14R Distributed by MAJOR® PHARMACEUTICALS, 17177 N. Laurel Park Drive, Suite 233, Livonia, MI 48152 Re-order No. 239567

6 0926127060 3



TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Rev. 06/17

Drug Facts

Active ingredient (in each tablet) Purpose

Bisacodyl 5 mg. Stimulant laxative

Uses

- relieves occasional constipation and irregularity
- this product generally produces bowel movement in 6 to 12 hours

Warnings

Do not use if you cannot swallow without chewing

Ask a doctor before use if you have

- stomach pain, nausea or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

When using this product

- do not chew or crush tablet(s)
- do not use within 1 hour after taking an antacid or milk
- you may have stomach discomfort, faintness or cramps

PEEL HERE FOR MORE DRUG FACTS

LOT #:

EXP. DATE: