MAJOR®

NDC 0904-7725-12

Loperamide Hydrochloride Tablets, 2 mg

Anti-Diarrheal

Controls the Symptoms of Diarrhea

12 CAPLETS*

COMPARE TO active ingredient of IMODIUM® A-D**

*Each Caplet (Capsule-Shaped Tablet) Contains 2 mg Loperamide Hydrochloride

**Major® Anti-Diarrheal is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Imodium® A-D.

FOR YOUR FAMILY'S PROTECTION, THIS PACKAGE IS CHILD RESISTANT. PLEASE FOLLOW INSTRUCTIONS ON THE BACK SIDE OF THE BLISTER CARD TO OPEN. IF DIFFICULT TO OPEN, USE SCISSORS.

CONVENIENT RECLOSING TAB

OPEN OTHER END

CODE AREA

12543 SC CV

Drug Facts

Active Ingredient (in each caplet)	Purpose
Loperamide HCl 2 mg	Anti-diarrheal

0904-7725-12 0

Drug Facts (continued)

Use controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Do not use if you have bloody or black stool

Ask a doctor before use if you have
- fever
- mucus in the stool
- a history of liver disease

Ask a doctor or pharmacist before use if you are taking antibiotics

When using this product
- dizziness, drowsiness or dizziness may occur. Be careful when driving or operating machinery

Stop use and ask a doctor if
- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away

Directions
- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool, 1 caplet after each subsequent loose stool, but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool, 1/2 caplet after each subsequent loose stool, but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool, 1/2 caplet after each subsequent loose stool, but no more than 2 caplets in 24 hours
children under 6 years (up to 47 lbs)	ask a doctor

Other information
- store between 20-25°C (68-77°F)
- **do not use if carton or blister unit is broken or torn**
- see end panel for lot number and expiration date

Drug Facts (continued)

Inactive ingredients anhydrous lactose, camauba wax, D&C yellow no. 10, FD&C blue no. 1, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch

Questions or comments?
1-800-719-9260

Distributed By: MAJOR PHARMACEUTICALS
37779 Enterprise Drive, LIVONIA, MI 48150, USA
M-06 REV 0908
Re-Order No. 700793

42355422

22435SCC4