

Drug Facts

Active ingredient (in each 15 mL Tbsp.) Purpose

Bismuth subsalicylate 262 mgUpset stomach reliever
and anti-diarrheal

Uses relieves: ■ diarrhea ■ heartburn ■ indigestion
■ nausea ■ upset stomach associated with these symptoms

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are

■ allergic to salicylates (including aspirin) ■ taking other salicylate products

Do not use if you have ■ bloody or black stool ■ an ulcer
■ a bleeding problem

Ask a doctor before use if you have ■ fever ■ mucus in the stool

Ask a doctor or pharmacist before use if you are taking
any drug for ■ anticoagulation (thinning of the blood)
■ diabetes ■ gout ■ arthritis

When using this product a temporary, but harmless darkening of the stool and/or tongue may occur.

Stop use and ask a doctor if ■ symptoms get worse
■ ringing in the ears or loss of hearing occurs
■ diarrhea lasts more than 2 days

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center immediately.

◀ **Drug Facts** continued

MFR# 53041

NDC 0904-1313-09

 **MAJOR®**

Bismatrol

**Soothing, coating action provides
fast relief for heartburn,
indigestion, nausea,
upset stomach and diarrhea**

***COMPARE TO THE ACTIVE
INGREDIENT IN PEPTO-BISMOL®.**

**8 FL OZ
(236 mL)**

Drug Facts (continued)

Directions ■ **shake well before use** ■ for accurate dosing, use dose cup ■ adults and children 12 years and over: 1 dose (2 Tbsp or 30 mL) every 1/2 to 1 hour as needed ■ do not exceed 8 doses (16 Tbsp or 240 mL) in 24 hours ■ use until diarrhea stops but not more than 2 days ■ children under 12 years: ask a doctor ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information ■ each Tbsp contains:

• sodium 5 mg • salicylate 130 mg • potassium 15 mg
■ protect from freezing. ■ avoid excessive heat (over 104°F or 40°C). ■ **TAMPER EVIDENT: Do not use if imprinted shrinkband is missing or broken.**

Inactive ingredients benzoic acid, D&C red 22, D&C red 28, flavor, hydroxyethyl cellulose, potassium hydroxide, purified water, saccharin sodium, salicylic acid, simethicone, xanthan gum.

Questions or Comments? (800) 616-2471

*This product is not manufactured or distributed by Procter & Gamble, Inc., the distributor of Pepto-Bismol®.

Distributed By MAJOR® PHARMACEUTICALS
31778 Enterprise Drive, Livonia, MI 48150 USA

Rev. 04/12

Re-Order No. 700281

M-30

