

COATING FREE

## Drug Facts

### Active ingredient (in each tablet)

Bismuth subsalicylate 262 mg .....Upset stomach reliever  
(total salicylate 102 mg per tablet) and anti-diarrheal

### Purpose

**Uses** relieves: ■ diarrhea ■ heartburn ■ indigestion ■ nausea  
■ upset stomach associated with these symptoms

### Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Contains salicylate. Do not take if you are ■ allergic to salicylates (including aspirin) ■ taking other salicylate products

**Do not use if you have** ■ bloody or black stool ■ an ulcer  
■ a bleeding problem

**Ask a doctor before use if you have** ■ fever ■ mucus in the stool

**Ask a doctor or pharmacist before use if you are** taking any drug for  
■ anticoagulation (thinning of the blood) ■ diabetes ■ gout ■ arthritis

**When using this product** a temporary, but harmless darkening of the stool and/or tongue may occur.

**Stop use and ask a doctor if** ■ symptoms get worse  
■ ringing in the ears or loss of hearing occurs  
■ diarrhea lasts more than 2 days

**If pregnant or breast-feeding, ask a health professional before use.**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center immediately.

**Directions** ■ chew or dissolve in mouth ■ adults and children 12 years and over: 2 tablets every 1/2 to 1 hour as needed ■ do not take more than 8 doses (16 tablets) in 24 hours ■ children under 12 years: ask a doctor  
■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

### Drug Facts (continued)

**Other information** ■ each tablet contains: ● sodium less than 1 mg ● salicylate 102 mg ● calcium 140 mg  
■ very low sodium ■ avoid excessive heat (over 104°F or 40°C). ■ **TAMPER EVIDENT: Do not use if individual compartments are torn or open.**

**Inactive ingredients** calcium carbonate, D&C red 27 aluminum lake, flavor, magnesium stearate, mannitol, pregelatinized starch, saccharin sodium.

# Bismatrol

## Chewable Tablets

Soothing, coating action provides fast relief for heartburn, indigestion, nausea, upset stomach and diarrhea



NDC 0904-1315-46

# Bismatrol

## Chewable Tablets

Soothing, coating action provides fast relief for heartburn, indigestion, nausea, upset stomach and diarrhea

\*COMPARE TO THE ACTIVE  
INGREDIENT IN PEPTO-BISMOL®.

30 CHEWABLE  
TABLETS



MFR# 53041 REV 0312

M234690



**Bismatrol**  
Chewable Tablets  
**30 CHEWABLE  
TABLETS**

\* This product is not manufactured or distributed by Procter & Gamble, Inc., owner of the registered trademark PEPTO-BISMOL®

Distributed By MAJOR PHARMACEUTICALS  
31778 Enterprise Drive, Livonia, MI 48150 USA

Re-Order No. 700286 M-30 Rev.03/12

COATING FREE