

FOR YOUR SAFETY, SUPPOSITORIES ARE
PACKAGED IN TAMPER RESISTANT
PLASTIC WRAPPER. DO NOT USE IF
PLASTIC WRAPPER IS TORN OR OPEN

brand of
HEMORRHOIDAL
SUPPOSITORIES

Ann-Med

12 SUPPOSITORIES

RELIEVES HEMORRHOIDAL PAIN AND BURNING

brand of
HEMORRHOIDAL
SUPPOSITORIES

Ann-Med



MAJOR®

NDC 0904-7688-22

HEMORRHOIDAL TISSUE

- NON-STAINING
- FAST ACTING
- PROTECTS IRRITATED TISSUE
- SHRINKS SWOLLEN

brand of
HEMORRHOIDAL
SUPPOSITORIES

Ann-Med

Drug Facts

Active ingredients (in each suppository) **Purposes**
Hard Fat 88.7%Protectant
Phenylephrine HCl 0.25%Vasoconstrictor

Uses

- temporarily shrinks hemorrhoidal tissue
- relieves the itching, burning and discomfort associated with hemorrhoids

Warnings

For rectal use only

Ask a doctor before use if you have

- heart disease ■ thyroid disease ■ high blood pressure
- diabetes ■ trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are presently taking a prescription drug for high blood pressure or depression

When using this product, do not use more than directed

Stop use and ask a doctor if

- bleeding occurs
- condition worsens or does not improve within 7 days

Drug Facts (continued)

If **pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- when practical, cleanse the affected area by patting or blotting with an appropriate cleansing wipe.
 - gently dry by patting or blotting with a tissue or a soft cloth before insertion of suppository.
 - detach one suppository from the strip
 - separate the two tabs of the white plastic wrap
 - with thumb and forefinger of each hand, grasp the tabs and peel downward, unwrapping the suppository
 - insert one suppository into the rectum
- adults and children 12 years and over:** insert one suppository rectally up to 4 times daily
- children under 12 years:** ask a doctor

Questions?

Serious Adverse Effects call: (800) 616-2471

Other information

store below 30°C (86°F) to avoid melting.

Inactive ingredients

corn starch, methylparaben, propylparaben

Distributed by:

MAJOR PHARMACEUTICALS
31178 Enterprise Drive
Livonia, MI 48150, U.S.A.

M-64

Re-Order No. 700419



6053-0610