

MAJOR

16 TABLETS

Cough & Cold

Relief for people with
High Blood Pressure 

NDC 0904-5817-44

**Compare to the active ingredients in Coricidin[®] HBP Cough & Cold*

MAJOR

Cough & Cold

Relief for people with
High Blood Pressure 

Chlorpheniramine maleate - Antihistamine
Dextromethorphan HBr - Cough Suppressant

Decongestant Free

Relieves

• Cough • Runny Nose • Sneezing

16 TABLETS



TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

No print/No varnish
Lot & Exp date

Drug Facts (continued)


Questions or comments? (800) 616-2471

Drug Facts (continued)

microcrystalline cellulose, povidone, propylene glycol, propylparaben, stearic acid, silica gel, sodium benzoate, stearic acid, sugar, talc, titanium dioxide

*This product is not manufactured or distributed by MSD Consumer Care, Inc., owner of the registered trademark Coricidin[®] HBP Cough & Cold.
50844 RE/0714D41121

Distributed by **MAJOR[®] PHARMACEUTICALS**
31778 Enterprise Drive, Livonia, MI 48150 USA M-17
Re-order No. 873602 Rev. 08/14



Drug Facts

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Active ingredients (in each tablet)

Chlorpheniramine maleate 4 mg. Antihistamine
Dextromethorphan HBr 30 mg. Cough suppressant

Purpose

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- temporarily relieves runny nose and sneezing due to the common cold

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- trouble urinating due to an enlarged prostate gland
- glaucoma
- cough that occurs with excessive phlegm (mucus)
- a breathing problem or persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.

When using this product

- excitability may occur, especially in children

Drug Facts (continued)

- marked drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if persistent cough lasts for more than one week, tends to recur, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

do not exceed recommended dose

- adults and children 12 years and over: 1 tablet every 6 hours. Do not take more than 4 tablets in 24 hours.
- children under 12 years: do not use

Other information

TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

acacia, black iron oxide, calcium carbonate, carnauba wax, croscarmellose sodium, D&C red #27 aluminum lake, FD&C yellow #6 aluminum lake, lactose, magnesium stearate, methylparaben,

01945r1

B-1212-411-21
REV0714D41121