



Day Time

MULTI-SYMPTOM
COLD/FLU RELIEF



NDC 0904-6607-44

Compare to the active ingredients in
Vicks® DayQuil® Cold & Flu LiquiCaps®

Day Time

MULTI-SYMPTOM COLD/FLU RELIEF

ACETAMINOPHEN - Pain Reliever/Fever Reducer
DEXTROMETHORPHAN HBr - Cough Suppressant
PHENYLEPHRINE HCl - Nasal Decongestant

NON-DROWSY

Headache, Fever, Sore Throat,
Nasal Congestion, Cough

Alcohol-Free
Antihistamine-Free



16 Softgels

Distributed by: **MAJOR® PHARMACEUTICALS**
17177 N Laurel Park Drive, Suite 233
Livonia, MI 48152, USA
Re-order No. 700832
M-17 Rev. 10/16

*This product is not manufactured or distributed
by Procter & Gamble, owner of the registered
trademark Vicks® DayQuil® Cold & Flu LiquiCaps®.
Product of China
Packaged and Quality Assured in the USA

No print/No varnish
Lot & Exp date

TAMPER EVIDENT: DO NOT USE IF
PACKAGE IS OPENED OR IF ALUSTER
UNIT IS TORN, BROKEN OR SHOWS
ANY SIGNS OF TAMPERING



3 0904660744 1

Drug Facts (continued)	Other information
<p>■ see end flap for expiration date and lot number</p> <p>■ protect from heat, humidity and light</p> <p>(59°-86°F)</p> <p>■ store at 25°C (77°F); excursions permitted between 15°-30°C</p> <p>OR BLISTER IS TORN OR BROKEN</p> <p>■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED</p>	
Drug Facts (continued)	Inactive ingredients
	yellow #6, gelatin, glycerin, lecithin, mineral oil, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol
	Questions or comments? (800) 616-2471
	50844 ORG021665921

B-1212-659-21
ORG021665921

Drug Facts	Active ingredients (in each liquid-filled capsule)
<p>KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION</p> <p>Purpose</p> <p>Acetaminophen 325 mg Pain reliever/fever reducer Dextromethorphan HBr 10 mg Cough suppressant Phenylephrine HCl 5 mg Nasal decongestant</p> <p>Uses</p> <p>■ temporarily relieves common cold and flu symptoms: ■ sore throat ■ nasal congestion ■ headache ■ fever ■ minor aches and pains ■ cough due to minor throat and bronchial irritation</p> <p>Warnings</p> <p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product</p> <p>Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p> <p>Do not use</p> <p>■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ if you have ever had an allergic reaction to this product or any of its ingredients</p>	
Drug Facts (continued)	Ask a doctor before use if you have
	■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
	■ liver disease ■ thyroid disease ■ diabetes
	■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema ■ heart disease ■ high blood pressure
	■ cough that occurs with too much phlegm (mucus)
	■ difficulty in urination due to enlargement of the prostate gland
	■ thinning drug warfarin.
	When using this product do not exceed recommended dosage.
	Stop use and ask a doctor if:
	■ nervousness, dizziness, or sleeplessness occur
	■ pain, nasal congestion, or cough gets worse or lasts more than 7 days
	■ fever gets worse or lasts more than 3 days
	■ redness or swelling is present ■ new symptoms occur
	■ accompanied by fever, rash, or persistent headache
	These could be signs of a serious condition.
	If pregnant or breast-feeding, ask a health professional before use.
	Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
	Directions ■ do not take more than directed
	■ adults and children 12 years and over: take 2 capsules with water every 4 hours
	■ children under 12 years: ask a doctor