



DO NOT USE IF PRINTED NECKBAND
IS BROKEN OR MISSING

ANTIHISTAMINE, Brompheniramine Maleate
NASAL DECONGESTANT, Phenylephrine HCl

DimaphenTM

Children's

MAJOR

NDC 0904-6462-09

MAJOR

Children's

DimaphenTM

ANTIHISTAMINE, Brompheniramine Maleate
NASAL DECONGESTANT, Phenylephrine HCl

NDC 0904-6462-09

MAJOR

Children's

DimaphenTM

ANTIHISTAMINE, Brompheniramine Maleate
NASAL DECONGESTANT, Phenylephrine HCl

Relieves • Nasal Symptoms
• Stuffy Nose • Runny Nose
• Sneezing
Plus Other Symptoms
• Itchy, Watery Eyes

GRAPE FLAVOR



Alcohol-
Free

6 yrs.
and older

Relieves • Nasal Symptoms
• Stuffy Nose • Runny Nose
• Sneezing
Plus Other Symptoms
• Itchy, Watery Eyes

GRAPE FLAVOR



Alcohol-
Free

6 yrs.
and older

COMPARE TO
the active ingredients of
DIMETAPP® COLD & ALLERGY*

8 FL. OZ. (237 mL.)

COMPARE TO
the active ingredients of
DIMETAPP® COLD & ALLERGY*

8 FL. OZ. (237 mL.)

Drug Facts

Active ingredients (in each 10 mL)

Brompheniramine maleate, USP 2 mg.....Antihistamine
Phenylephrine HCl, USP 5 mg.....Nasal decongestant

Purposes

Uses

- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- temporarily relieves these symptoms due to hay fever (allergic rhinitis):
 - runny nose
 - sneezing
 - itchy, watery eyes
 - itching of the nose or throat
- temporarily restores freer breathing through the nose

Warnings

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- to make a child sleepy

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product

- do not use more than directed
- drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- you get nervous, dizzy, or sleepless
- symptoms do not get better within 7 days or are accompanied by fever

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Drug Facts (continued)

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosage cup provided
- keep dosage cup with product
- mL = milliliter

age	dose
adults and children 12 years and over	20 mL every 4 hours
children 6 to under 12 years	10 mL every 4 hours
children under 6 years	do not use

Other information

- each 10 mL contains: sodium 4 mg
- store at 20-25°C (68-77°F)

Inactive ingredients anhydrous citric acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sorbitol solution

Questions or comments? 1-800-616-2471

*Major® Dimaphen™ is not manufactured or distributed by Pfizer, distributor of Dimetapp® Cold & Allergy.

Gluten Free

Distributed by
MAJOR®
PHARMACEUTICALS
31778 Enterprise Drive
Livonia, MI 48150 USA
M-05 REV. 02/16
Re-Order No. 700411



3 09046 46209 5

LOT NO.

EXP.

CODE
AREA

: 90634 5C C5

