

Drug Facts

Active ingredients (in each 118 mL delivered dose)

| | |
|---------------------------------|-----------------|
| Dibasic sodium phosphate 7 g | Saline laxative |
| Monobasic sodium phosphate 19 g | Saline laxative |

Purpose

Uses

- relieves occasional constipation

Warnings

For rectal use only

Dosage warning: Using more than one enema in 24 hours can be harmful.

Ask a doctor before use if you have

- a sodium-restricted diet
- abdominal pain, nausea or vomiting
- kidney disease or are dehydrated
- noticed a sudden change in bowel habits that lasts over 2 weeks

Ask a doctor or pharmacist before use if you are taking any other drug. Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

When using this product do not use

- more than directed. Serious side effects may occur from excess dosage.
- for more than 3 days

Stop use and ask a doctor if

- you have rectal bleeding
- you have no bowel movement after use of a laxative

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

Do not use more unless directed by a doctor. See Warnings.

| | |
|--|-----------------------|
| adults & children 12 years and over | one bottle once daily |
| children 2 to under 12 years | 1/2 bottle once daily |
| children under 2 years | do not use |

Other information

- each 118 mL contains: sodium 4.3 g
- store at room temperature 20°-25°C (68°-77°F)
- this product generally produces bowel movement in 1 to 5 minutes

Inactive ingredients

benzalkonium chloride, disodium EDTA, purified water

Questions or comments?

1-800-616-2471


Directions

CAUTION: Remove blue protective shield before inserting: hold bottle upright, grasping bottle cap with fingers. Grasp protective shield with other hand and pull gently to remove.



Positioning:

Left-side position: lie on left side with knee bent and arms at rest.



Knee-chest position: kneel, then lower head and chest forward until left side of face is resting on surface. Position arms comfortably.



Administering enema:

- with steady pressure, gently insert enema into rectum with tip pointing toward navel
- squeeze bottle until recommended dose is expelled (it is not necessary to empty unit completely. Bottle contains more liquid than needed for effective use. A small amount of liquid will remain in bottle after squeezing).
- remove tip from rectum
- stop using if tip is hard to insert. Forcing the tip into the rectum can cause injury (especially if you have hemorrhoids). If enema tip causes rectal bleeding or pain, get immediate medical care.
- maintain position until urge to evacuate is strong (usually 1 to 5 minutes)
- if no urge is felt after 5 minutes of using, try to empty bowel. Call a doctor promptly if no liquid comes out of the rectum after 30 minutes because dehydration could occur.

Distributed by
MAJOR® PHARMACEUTICALS
31778 Enterprise Drive
LIVONIA, MI 48150, USA
M-139 Rev. 03/13
Re-Order No. 700202



3 09046 32078 4

Ready-to-Use
ENEMA
SALINE LAXATIVE

MAJOR

COMPARE TO the active
ingredients of FLEET®
ENEMA SALINE*

NDC 0904-6320-78

MAJOR

Ready-to-Use
ENEMA
SALINE LAXATIVE

For relief of
occasional
constipation
Pre-mixed

Soft, Flexible
Comfort Tip



LATEX FREE

4.5 FL OZ (133 mL)

COMPARE TO active
ingredients of FLEET®
ENEMA SALINE*

MAJOR

Ready-to-Use
ENEMA
SALINE LAXATIVE

For relief of occasional
constipation
Pre-mixed

Soft, Flexible
Comfort Tip

Protective shield
prevents
contamination

Comfortable,
lubricated tip for
ease of insertion

One-way safety valve
controls flow and
prevents reflux

Easy, soft squeeze
bottle



LATEX FREE

4.5 FL OZ (133 mL)