

Drug Facts (continued)

Directions

- Take with 8 to 12 glasses of water.
- Do not exceed 6 doses in 24 hours.

Storage

- Store at 25°C (77°F); excursions permitted between 15-30°C (59-86°F).

children under 6 years of age consult a physician

adults and children 10 years of age

children 6 to 12 years 1/2 to 1 tablet every 4 hours

adults and children 10 years of age 4 hours

children under 6 years of age consult a physician

adults and children 10 years of age 1 to 2 tablets every 4 hours

children 6 to 12 years 1/2 to 1 tablet every 4 hours

adults and children 10 years of age 4 hours

children under 6 years of age consult a physician

adults and children 10 years of age 1 to 2 tablets every 4 hours

children 6 to 12 years 1/2 to 1 tablet every 4 hours

adults and children 10 years of age 4 hours

children under 6 years of age consult a physician

adults and children 10 years of age 1 to 2 tablets every 4 hours

children 6 to 12 years 1/2 to 1 tablet every 4 hours

adults and children 10 years of age 4 hours

children under 6 years of age consult a physician

adults and children 10 years of age 1 to 2 tablets every 4 hours

children 6 to 12 years 1/2 to 1 tablet every 4 hours

adults and children 10 years of age 4 hours

children under 6 years of age consult a physician

adults and children 10 years of age 1 to 2 tablets every 4 hours

children 6 to 12 years 1/2 to 1 tablet every 4 hours

adults and children 10 years of age 4 hours

children under 6 years of age consult a physician

adults and children 10 years of age 1 to 2 tablets every 4 hours

children 6 to 12 years 1/2 to 1 tablet every 4 hours

adults and children 10 years of age 4 hours

children under 6 years of age consult a physician

adults and children 10 years of age 1 to 2 tablets every 4 hours

children 6 to 12 years 1/2 to 1 tablet every 4 hours

adults and children 10 years of age 4 hours

NDC 0904-5154-60



GUAIIFENESIN TABLETS

Guaiifenesin 200 mg

Expectorant

- Relieves Chest Congestion
- Thins and Loosens Mucus



100 Tablets

Immediate Release

REV0812B8812

Distributed by MAJOR® PHARMACEUTICALS

31778 Enterprise Drive, Livonia, MI 48150 USA
Re-Order No. 238163 M-17 Rev. 04/13



TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts

Purpose (in each immediate-release tablet)

Expectorant
Guaiifenesin 200 mg

Uses
Helps clear mucus from the bronchial passageways of both lungs and makes coughs more productive.

Warnings

Ask a doctor before use if you have

- sinusitis or chronic bronchitis such as occurs with sinusitis, asthma, chronic bronchitis, or emphysema
 - colds accompanied by too much phlegm (mucus)
- Stop use and ask a doctor if a cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious illness.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

PEEL HERE FOR MORE DRUG FACTS

STOP PEELING

Question or comment?

1-774-961-0010 (100)