

Drug Facts**Active ingredient (in each 7.5 mL) Purpose**Loperamide HCl
1 mg.....Anti-diarrheal**Use** controls symptoms of diarrhea, including Travelers' Diarrhea**Warnings****Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl**Do not use** if you have bloody or black stool**PEEL BACK HERE** **MAJOR**® NDC 0904-6256-20**Loperamide Hydrochloride Oral Suspension**

Anti-Diarrheal Liquid

Controls the Symptoms of Diarrhea

Mint Flavor

4 FL OZ (120 mL)

1 mg Loperamide Hydrochloride per 7.5 mL

COMPARE TO the active ingredient of IMODIUM® A-D***Do not use if printed plastic neckband is broken or missing**

*Major® Loperamide Hydrochloride Oral Suspension is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Imodium® A-D.

Distributed by: **MAJOR® PHARMACEUTICALS**
31778 Enterprise Drive, Livonia, MI 48150 USA
IM-06 REV. 09/13
Re-Order No. 700606**Gluten Free****CODE AREA**• no varnish • no color
: 64526 5C F2**ADHESIVE AREA NO COPY****Drug Facts (continued)**

children 6-8 years (48-59 lbs)

15 mL (3 tsp) after the first loose stool; 7.5 mL (1 1/2 tsp) after each subsequent loose stool; but no more than 30 mL (6 tsp) in 24 hours

children under 6 years (up to 47 lbs)

ask a doctor

Other information■ each 30 mL (6 tsp) contains: sodium 15 mg
■ store between 20-25°C (68-77°F)**Drug Facts (continued)**

■ see side panel for lot number and expiration date

Inactive ingredients anhydrous citric acid, carboxymethylcellulose sodium, D&C yellow no. 10, FD&C blue no. 1, glycerin, microcrystalline cellulose, natural and artificial mint flavor, propylene glycol, purified water, simethicone, sodium benzoate, sucralose, titanium dioxide, xanthan gum**Questions or comments?**

1-800-719-9260

Drug Facts (continued)**Ask a doctor before use if you have** ■ fever
■ mucus in the stool ■ a history of liver disease**Ask a doctor or pharmacist before use if you are taking antibiotics****When using this product** ■ tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.**Stop use and ask a doctor if**■ symptoms get worse
■ diarrhea lasts for more than 2 days
■ you get abdominal swelling or bulging.
These may be signs of a serious condition.**If pregnant or breast-feeding**, ask a health professional before use.**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison**Drug Facts (continued)**

Control Center right away. (1-800-222-1222)

Directions ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea
■ find right dose on chart. If possible, use weight to dose; otherwise use age. ■ shake well before using ■ only use attached measuring cup to dose product

adults and children 12 years and over

30 mL (6 tsp) after the first loose stool; 15 mL (3 tsp) after each subsequent loose stool; but no more than 60 mL (12 tsp) in 24 hours

children 9-11 years (60-95 lbs)

15 mL (3 tsp) after the first loose stool; 7.5 mL (1 1/2 tsp) after each subsequent loose stool; but no more than 45 mL (9 tsp) in 24 hours

ADHESIVE AREA NO COPY