

Rugby[®]

Compare to
MiraLAX[®]*

NDC 0536-1052-27

Original Prescription Strength

PEG3350

*Polyethylene Glycol 3350 Powder
for Oral Solution, Osmotic Laxative*

- **Relieves Occasional Constipation (irregularity)**
- **Softens Stool**

NET WT 17.9 OZ (510 gm)

30 ONCE-DAILY DOSES

.5625" White Space Area

*This product is not manufactured or distributed by MSD Consumer Care, Inc., the distributor of MiraLAX[®].

Distributed by:
RUGBY LABORATORIES
17177 N Laurel Park Drive, Suite 233
Livonia, MI 48152
www.rugbylaboratories.com

R-132 Rev. 08/16
Re-Order No. 370393

LA0522702902



Unvarnished area

.5625" White Space Area

Drug Facts

Active ingredient (in each dose) Purpose

Polyethylene Glycol 3350,
17 g (cap filled to line)..... Osmotic Laxative

Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

Warnings

Allergy alert: Do not use if you are allergic to polyethylene glycol

Do not use if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor or pharmacist before use if you are taking a prescription drug

When using this product you may have loose, watery, more frequent stools

Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a POISON CONTROL CENTER right away. (1-800-222-1222)

Directions

- **do not take more than directed unless advised by your doctor**
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line
- adults and children 17 years of age and older:
 - fill to top of line in cap which is marked to indicate the correct dose (17 g)
 - stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
 - use once a day
 - use no more than 7 days
- Children 16 years of age or under: ask a doctor

Other information

- store at 20° - 25°C (68° - 77°F)
- tamper-evident: do not use if printed foil seal under cap is missing, open or broken

Inactive ingredients

none

Questions or comments?

1-800-645-2158

Polyethylene Glycol 3350 increases frequency of bowel movements and softens the stool

*Dissolves in Any Beverage
Sugar free*