



Omeprazole

Delayed Release Tablets 20 mg
Acid Reducer

Drug Facts (continued)

Active ingredient (in each tablet) Omeprazole 20 mg.....acid reducer

Use
 ■ treats frequent heartburn (occurs 2 or more days a week)
 ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Allergy alert: Do not use if you are allergic to omeprazole with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
Do not use if you have trouble or pain swallowing food, vomiting or chest pain or shoulder pain with shortness of breath, sweating, heartburn with lightheadedness, sweating or dizziness, chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness, frequent wheezing, particularly with heartburn, frequent weight loss, unexplained weight loss, nausea or vomiting, stomach pain

Ask a doctor or pharmacist before use if you are taking
 ■ warfarin, clopidogrel or clobazam (blood-thinning medicines)
 ■ prescription antifungal or anti-yeast medicines
 ■ dexamethasone (steroid medicine)
 ■ digoxin (heart medicine)
 ■ tacrolimus or mycophenolate mofetil (immune system medicines)
 ■ prescription antivirals (medicines for HIV infection)
 ■ methotrexate (arthritis medicine)

Drug Facts (continued)

Other information
 ■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.
 ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor

Repeated 14-Day Courses (if needed)
 ■ you may repeat a 14-day course every 4 months

14-Day Course of Treatment
 ■ take every day for 14 days
 ■ do not take more than 1 tablet a day
 ■ do not use for more than 14 days unless directed by your doctor
 ■ swallow whole. Do not chew or crush tablets

Directions
 ■ for adults 18 years of age and older
 ■ this product is to be used once a day (every 24 hours), every day for 14 days
 ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

If pregnant or breast-feeding, ask a health professional before use:
 ■ you get diarrhea
 ■ you need to take more than 1 course of treatment every 4 months
 ■ your heartburn continues or worsens

Stop use and ask a doctor if
 ■ your heartburn continues or worsens

14-Day Course of Treatment
 ■ take every day for 14 days
 ■ do not take more than 1 tablet a day
 ■ do not use for more than 14 days unless directed by your doctor
 ■ swallow whole. Do not chew or crush tablets

Repeated 14-Day Courses (if needed)
 ■ you may repeat a 14-day course every 4 months

Do not take for more than 14 days or more often than every 4 months unless directed by a doctor

Other information
 ■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.
 ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor

Other information
 ■ read the directions and warnings before use
 ■ keep the carton. It contains important information.
 ■ store at 20-25°C (68-77°F) and protect from moisture

Questions or comments? 1-800-719-9260

Questions or comments? 1-800-719-9260

Drug Facts (continued)

Inactive ingredients
 carnauba wax, ferric oxide red, ferric oxide yellow, hypromellose, hypromellose acetate succinate, lactose monohydrate, titanium dioxide, triethyl citrate

Tips for Managing Heartburn
 ■ Do not lie flat or bend over after eating
 ■ Do not wear tight-fitting clothing around the stomach
 ■ Do not eat before bedtime
 ■ Raise the head of your bed
 ■ Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 ■ Eat slowly and avoid big meals
 ■ If overweight, lose weight
 ■ Quit smoking

Made in Israel
Distributed by: MAJOR® PHARMACEUTICALS
31778 Enterprise Drive, Livonia, MI 48150 USA
M-05 REV. 04/15
Re-Order No. 263644



CODE AREA

915555CC9

ement of UPC is to be changed. is for customers ho are taking o shrink wraps.



Omeprazole

Delayed Release Tablets 20 mg
Acid Reducer

Safety Feature - Do not use if printed tablet blister unit is open or torn.



NDC 0904-5834-42

Omeprazole

Delayed Release Tablets 20 mg
Acid Reducer

Treats Frequent Heartburn!

42 Tablets
Three 14-day courses of treatment

Occurring 2 Or More Days A Week

Actual Size



EFB 1394
(915) 915-008 EFB1394-1

60055516

915555CC9