

No print/No varnish
Lot & Exp date



TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY
SEAL UNDER CAP IS BROKEN OR MISSING

B-1212-159B-12
REV0117E15912

Drug Facts (continued)

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take more or for a longer time than directed
- Caffeine warning:** The recommended dose of this product contains about as much **caffeine** as a cup of coffee. Limit the use of **caffeine**-containing medications, foods, or beverages while taking this product because too much **caffeine** may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.
- Do not use**
 - if you have ever had an allergic reaction to acetaminophen, aspirin, or any other pain reliever/fever reducer
 - with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- Ask a doctor before use if**
 - you have liver disease
 - stomach bleeding warning applies to you
 - you have a history of stomach problems, such as heartburn
 - you have high blood pressure, heart disease, liver cirrhosis or kidney disease
 - you have asthma
 - you are taking a diuretic
- Ask a doctor or pharmacist before use if you are**
 - taking a prescription drug for diabetes, gout, or arthritis
 - taking any other drug or are under a doctor's care for any serious condition

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- an allergic reaction occurs. Seek medical help right away, if any new symptoms appear
- ringing in the ears or loss of hearing occurs
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- painful area is red or swollen.

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not use more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over
 - take 2 tablets every 6 hours
 - do not take more than 8 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

corn starch, croscollon, hypromellose, microcrystalline cellulose, povidone, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments?

(800) 616-2471

*This product is not manufactured or distributed by Novartis AG, owner of the registered trademark Excedrin® Extra Strength.

50844 REV0117E15912

Distributed by MAJOR PHARMACEUTICALS

17171 W Laurel Park Drive, Suite 233
Livonia, MI 48152 USA M-17 Rev. 03/17
Re-order No. 252831

Drug Facts

Active ingredients (in each tablet)

Acetaminophen 250 mg Pain reliever
Aspirin 250 mg (NSAID)* Pain reliever
Caffeine 65 mg Pain reliever aid
*nonsteroidal anti-inflammatory drug

Uses ■ temporarily relieves minor aches and pains due to:

- headache
- arthritis
- a cold
- muscular aches
- toothache
- premenstrual and menstrual cramps

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- shock
- facial swelling
- hives
- asthma (wheezing)

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount
- 3 or more alcoholic drinks every day while using this product

■ with other drugs containing acetaminophen

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older

■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

■ have 3 or more alcoholic drinks every day while using this product

MAJOR

PAIN RELIEVER PLUS Acetaminophen, Aspirin (NSAID) and Caffeine

Pain Reliever / Pain Reliever Aid

100 COATED TABLETS

COMPARE TO the active ingredients in EXCEDRIN® Extra Strength†

NDC 0904-5135-59

MAJOR

PAIN RELIEVER PLUS Acetaminophen, Aspirin (NSAID) and Caffeine

Pain Reliever / Pain Reliever Aid

100 COATED TABLETS

COMPARE TO the active ingredients in EXCEDRIN® Extra Strength†

DC1193BR2