


3 0904-5529-87 4



Other information ■ read the directions and warnings before use

■ children under 12 years: ask a doctor
 ■ do not use more than 2 tablets in 24 hours
 ■ or drinking beverages that cause heartburn
 ■ to prevent symptoms, swallow 1 tablet with a glass of water. Do not chew.
 ■ to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.
 ■ adults and children 12 years and over.

Directions

Center right away.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

If pregnant or breast-feeding, ask a health professional before use.

■ you need to take this product for more than 14 days
 ■ your heartburn continues or worsens

Stop use and ask a doctor if

■ unexplained weight loss ■ nausea or vomiting ■ stomach pain
 ■ or lightheadedness
 ■ frequent chest pain ■ chest pain or shoulder pain with shortness of breath, sweating, or dizziness
 ■ heartburn with frequent wheezing, particularly with heartburn.
 ■ Ask a doctor before use if you have ■ had heartburn over 3 months. This may be a sign of a more serious condition.
 ■ These may be signs of a serious condition. See your doctor. ■ with other acid reducers

Do not use ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools.
 ■ Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Warnings

■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Uses

■ relieves heartburn associated with acid indigestion and sour stomach
 ■ famotidine 10 mg.

Active ingredient (in each tablet)

Purpose

Acid reducer

Drug Facts

DO NOT USE IF BLISTER UNIT IS BROKEN OR TORN

MAJOR® 

Heartburn Relief

Famotidine Tablets, 10 mg
Acid Reducer

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

MAJOR® 

NDC 0904-5529-87

Easy Push-Thru Package

Original Strength

Heartburn Relief

Famotidine Tablets, 10 mg
Acid Reducer
Just One Tablet

Prevents & Relieves Heartburn due to Acid Indigestion

COMPARE TO the active ingredient
of ORIGINAL STRENGTH PEPcid® AC*

ACTUAL SIZE



30 Tablets

Tablets

Drug Facts (continued)

■ keep the carton. It contains important information.
 ■ store at 20°-25°C (68°-77°F) ■ protect from moisture and light

Inactive ingredients carnauba wax, hypromellose, iron oxides, magnesium stearate, microcrystalline cellulose, pregelatinized starch, polyethylene glycol, talc, titanium dioxide, triacetin

(Questions or comments? 1-800-719-9260)

*Major® Heartburn Relief is not manufactured or distributed by Johnson & Johnson • MERCK Consumer Pharmaceutical Co., distributor of Original Strength PEPcid® AC.

Distributed By MAJOR PHARMACEUTICALS
31778 Enterprise Drive,
LIVONIA, MI 48150, USA
M-05 REV. 01/09
Re-Order No. 248045

CONVENIENT RECLOSING TAB

Code Area

OPEN OTHER END