

Drug Facts (continued)

Directions

- do not take more than 6 doses in any 24-hour period
- use enclosed dosing cup only. Do not use any other device.
- this adult product is not intended for use in children under 12 years of age ■ TSP= teaspoonful

adults and children 12 years and over

2 TSP every 4 hours

children under 12 years

do not use

Other information

■ each TSP (5 mL) contains:
sodium 3 mg ■ store at 20°-25°C (68°-77°F). Do not refrigerate

TAMPER-EVIDENT: Do not use this product if inner foil seal over the mouth of the bottle is cut, torn, broken or missing

Inactive ingredients

artificial cherry flavor, citric acid, corn syrup, FD&C Red #40, glycerin, menthol, purified water, saccharin sodium, sodium benzoate

Questions or comments ? (800)-616-2471

Distributed by:

MAJOR® PHARMACEUTICALS

17177 N Laurel Park Drive,
Suite 233, Livonia, MI 48152

M-64 REV 11/16

Re-Order No. 263913



Lot / Exp.:

* This product is not manufactured or distributed by Wyeth Consumer Healthcare, distributor of Robitussin® DM. Robitussin® DM is a registered trademark of A. H. Robins Company, Incorporated.

NDC 0904-0053-00
TAMPER-EVIDENT
MAJOR®
PHARMACEUTICALS

ROBAFEN™ DM

cough formula

Cough Suppressant
Expectorant

*Compare to active ingredients of
Robitussin®Cough+Chest Congestion DM

Dextromethorphan HBr, USP 20 mg and
Guaifenesin, USP 200 mg per 10 mL

• **Alcohol free**

• **Non-narcotic**

4 FL. OZ. (118 mL)

Drug Facts

Active ingredients (in each 2 TSP (10 mL))

Dextromethorphan HBr, USP 20 mg.....Cough suppressant

Guaifenesin, USP 200 mg.....Expectorant

Uses

■ temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold ■ helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes.

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have ■ cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema ■ cough that occurs with too much phlegm (mucus).

Stop use and ask a doctor if cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

PEEL HERE