

Adhesive Area

Drug Facts (continued)

reliever/fever reducer

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma ■ you have not been drinking fluids ■ you have lost a lot of fluid due to vomiting or diarrhea

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout, or arthritis
- taking any other drug
- under a doctor's care for any serious condition

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
 - you experience any of the following signs of stomach bleeding: ■ feel faint ■ have bloody or black stools ■ vomit blood ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days
 - redness or swelling is present ■ new symptoms occur ■ ringing in the ears or a loss of hearing occurs
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional

before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

Other information

- store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients

anhydrous lactose, carmelum wax, colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10 aluminum lake, iron oxide ochre, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polyacrylate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate

Questions or comments? 1-800-616-2471

STOP PEEING

NDC 0904-7704-70

MAJOR[®]
ASPIRIN[®]
LOW[™]

LOW STRENGTH 81mg ASPIRIN
(NSAID)

Enteric Coated

1 1/4 gr. (81 mg) each Analgesic

250 Tablets

Compare to the active ingredient in
BAYER[®] Low Dose[™]

*This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Bayer[®] Low Dose[™]. 481R, Rev. 06/17, M-14

Distributed by

MAJOR[®] PHARMACEUTICALS

17177 N. Laurel Park Drive

Suite 233

Livonia, MI 48152

Re-order No. 009524



TAMPER EVIDENT DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts

Active ingredient (in each tablet)

1 ASPIRIN (NSAID) 81 mg Pain reliever
*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

■ temporarily relieves minor aches and pains
Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief. ■ ask your doctor about other uses for enteric-coated 81 mg Aspirin

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include: ■ hives ■ facial swelling ■ shock ■ asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Do not use if you are allergic to aspirin or any other pain

PEEL HERE
FOR MORE
DRUG FACTS

LOT #:
Varnish
Omit Area
EXP. DATE: