

ADHESIVE AREA NO COPY

■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for diabetes, gout or arthritis

Stop use and ask a doctor if ■ an allergic reaction occurs. Seek medical help right away.

■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood

■ have bloody or black stools

■ have stomach pain that does not get better ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts for more than 3 days ■ redness or swelling is present

■ new symptoms occur ■ ringing in the ears or loss of hearing occurs

These could be signs of a serious condition

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last

3 months of pregnancy unless definitely directed to do so by a doctor because it may

cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions ■ drink a full glass of water with each dose

■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor

■ children under 12 years: consult a doctor

Store at 20-25°C (68-77°F).

Contains FD&C yellow no. 6 aluminum lake

Questions or comments?

1-800-616-2471

**Talk to your doctor before you begin an aspirin regimen