


(604)
EFB 667
604-008 EFB667-1



Drug Facts

Active ingredient (in each tablet) **Purposes**
Ibuprofen 200 mg (NSAID)*Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - headache ■ muscular aches ■ minor pain of arthritis
 - toothache ■ backache ■ the common cold
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives ■ facial swelling ■ asthma (wheezing)
- shock ■ skin reddening ■ rash ■ blisters

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chances are higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit or aspirin
- under a doctor's care for any serious condition
- taking any other drug

NDC 0904-7915-70

MAJOR

Ibuprofen

Tablets, 200mg

PAIN RELIEVER / FEVER REDUCER (NSAID)

COMPARE TO the active ingredient of **ADVIL**®**

250 COATED TABLETS

Actual Size

SEE NEW WARNINGS INFORMATION

NDC 0904-7915-70

MAJOR

Ibuprofen

Tablets, 200mg

PAIN RELIEVER / FEVER REDUCER (NSAID)

COMPARE TO the active ingredient of **ADVIL**®**

250 COATED TABLETS

Actual Size

SEE NEW WARNINGS INFORMATION

Drug Facts (continued)

Questions or comments? 1-800-719-9260

KEEP CARTON FOR REFERENCE

DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING

**Major® Ibuprofen Tablets, 200mg are not manufactured or distributed by Wyeth Consumer Healthcare, distributor of Advil®.

Gluten Free

Distributed by: MAJOR PHARMACEUTICALS
31778 Enterprise Drive, Livonia, MI 48150 USA
M-05 REV. 07/11
Re-Order No. 700389

Drug Facts (continued)

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint ■ vomit blood ■ have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
(1-800-222-1222)

Directions


- **do not take more than directed**
- **the smallest effective dose should be used**

adults and children 12 years and older	■ take 1 tablet every 4 to 6 hours while symptoms persist
	■ if pain or fever does not respond to 1 tablet, 2 tablets may be used
	■ do not exceed 6 tablets in 24 hours, unless directed by a doctor
children under 12 years	■ ask a doctor

Other information

- read all warnings and directions before use
- store at 20-25°C (68-77°F)
- avoid high humidity and excessive heat above 40°C (104°F)
- see end panel for lot number and expiration date

Inactive ingredients
colloidal silicon dioxide, corn starch, croscarmellose sodium, hypromellose, iron oxide red, iron oxide yellow, microcrystalline cellulose, polyethylene glycol, polysorbate 80, stearic acid, titanium dioxide



LOT NO.

EXP.

CODE AREA

3 09047 91570 5