

Drug Facts (continued)

■ find the right dose on chart below. If possible, use weight to dose; otherwise, use age.

| Weight (lb) | Age (year) | Dose (mL)* |
|-------------|------------|--------------|
| under 24 | under 2 | ask a doctor |
| 24-35 | 2-3 | 5 mL |
| 36-47 | 4-5 | 7.5 mL |
| 48-59 | 6-8 | 10 mL |
| 60-71 | 9-10 | 12.5 mL |
| 72-95 | 11 | 15 mL |

*or as directed by a doctor

Other information ■ each 5 mL contains: sodium 2 mg ■ store between 20-25°C (68-77°F) ■ do not refrigerate. **Inactive ingredients** acesulfame potassium, butylparaben, carboxymethylcellulose sodium, citric acid, FD&C red #40, flavors, glycerin, high fructose corn syrup, microcrystalline cellulose, propylene glycol, purified water, sodium benzoate, sorbitol, xanthan gum

Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

Distributed by:
Major® Pharmaceuticals
 17177 N Laurel Park Drive, Suite 233
 Livonia, MI 48152

Re-Order No. 700530
 Rev. 09/16 M-58
 PLD-C231D LB004379



NDC 0904-6536-20

Children's
Māpap
 Acetaminophen
 160 mg per 5 mL

Oral Suspension

Pain reliever/Fever reducer

- Alcohol free
- Aspirin free
- Ibuprofen free

Cherry
 Flavor



For ages 2-11 years

4 fl oz (118 mL)

Drug Facts

Active ingredient (in each 5 mL)

Purpose

Acetaminophen 160 mg.....Pain reliever/fever reducer

Uses ■ temporarily reduces fever ■ temporarily relieves minor aches and pains due to ■ the common cold ■ flu ■ headache ■ sore throat ■ toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes: ■ more than 5 doses in 24 hours, which is the maximum daily amount

■ with other drugs containing acetaminophen. **Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away. **Sore throat**

warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. **Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription).

If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ if your child is allergic to acetaminophen or any of the inactive ingredients in this product. **Ask a doctor before use if your child has liver disease. Ask a doctor or**

pharmacist before use if your child is taking the blood thinning drug warfarin. When using this product, do not exceed recommended dose (see Overdose warning). Stop use and ask a doctor if ■ new symptoms occur ■ redness or swelling is present

■ pain gets worse or lasts more than 5 days ■ fever gets worse or lasts more than 3 days. These could be signs of a serious condition. **Keep out of reach of children.**

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms. **Directions** ■ this product does not contain directions or complete warnings for adult use ■ do not take more than directed (see Overdose warning) ■ shake well before using ■ mL = milliliter ■ use only the enclosed dosing cup designed for use with this product. Do not use any other dosing device.

■ if needed, repeat dose every 4 hours while symptoms last ■ do not give more than 5 times in 24 hours ■ do not give more than 5 days unless directed by a doctor

■ if needed, repeat dose every 4 hours while symptoms last ■ do not give more than 5 times in 24 hours ■ do not give more than 5 days unless directed by a doctor

■ if needed, repeat dose every 4 hours while symptoms last ■ do not give more than 5 times in 24 hours ■ do not give more than 5 days unless directed by a doctor

■ if needed, repeat dose every 4 hours while symptoms last ■ do not give more than 5 times in 24 hours ■ do not give more than 5 days unless directed by a doctor

■ if needed, repeat dose every 4 hours while symptoms last ■ do not give more than 5 times in 24 hours ■ do not give more than 5 days unless directed by a doctor