


833550250



**MAJOR**

For Ages 6 mos. to 23 mos.

**Infants'**  
**CONCENTRATED DROPS**

**Ibuprofen Oral**  
**Suspension**

50 mg per 1.25 mL

Pain Reliever/Fever Reducer  
(NSAID)


**Infants'**



Use only with  
enclosed syringe  
Berry Flavored  
Lasts up to 8 hours  
No High Fructose  
Corn Syrup

0.5 FL OZ (15 mL)

COMPARE TO  
the Active  
Ingredient of  
**MOTRIN®**  
**INFANTS' DROPS™**



057055CC8

NDC 0904-5663-35

Important: Read all product information before using. Keep this box for important information. This product is intended for use in children ages 6 months to 23 months of age.


**Drug Facts**

**Active ingredient** (in each 1.25 mL) **Purposes**  
Ibuprofen 50 mg.....Pain reliever/  
(NSAID)\* fever reducer  
\*nonsteroidal anti-inflammatory drug

**Uses** temporarily: ■ reduces fever  
■ relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

**Warnings**  
**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters  
If an allergic reaction occurs, stop use and seek medical help right away.  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child: ■ has had stomach ulcers or bleeding problems ■ takes a blood thinning (anticoagulant) or steroid drug ■ takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ takes more or for a longer time than directed

**Sore throat warning:** Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.



**Infants'**  
**CONCENTRATED DROPS**

**Ibuprofen Oral**  
**Suspension**

50 mg per 1.25 mL

Pain Reliever/Fever Reducer (NSAID)

Child Resistant Safety Cap • Read Instructions Carefully

**Drug Facts (continued)**

**Do not use** ■ if the child has ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery

**Ask a doctor before use if**  
■ stomach bleeding warning applies to your child  
■ child has a history of stomach problems, such as heartburn  
■ child has problems or serious side effects from taking pain relievers or fever reducers ■ child has not been drinking fluids  
■ child has lost a lot of fluid due to vomiting or diarrhea  
■ child has high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ child has asthma ■ child is taking a diuretic

**Ask a doctor or pharmacist before use if the child is**  
■ under a doctor's care for any serious condition ■ taking any other drug

**When using this product** ■ give with food or milk if stomach upset occurs ■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

**Stop use and ask a doctor if**  
■ child experiences any of the following signs of stomach bleeding:  
■ feels faint ■ vomits blood ■ has bloody or black stools  
■ has stomach pain that does not get better  
■ the child does not get any relief within first day (24 hours) of treatment  
■ fever or pain gets worse or lasts more than 3 days  
■ redness or swelling is present in the painful area  
■ any new symptoms appear

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions** ■ this product does not contain directions or complete warnings for adult use

■ **do not give more than directed** ■ shake well before using  
■ find right dose on chart below. If possible, use weight to dose; otherwise use age.  
■ measure with the dosing device provided. Do not use with any other device.  
■ dispense liquid slowly into the child's mouth, toward the inner cheek  
■ if needed, repeat dose every 6-8 hours  
■ do not use more than 4 times a day

**Dosing Chart**

Weight (lbs)	Age (mos)	Dose (mL)
under 6 mos		ask a doctor
12-17 lbs	6-11 mos	1.25 mL
18-23 lbs	12-23 mos	1.875 mL

**EFB 294**  
(057) 057-006 EFB294-1

**Drug Facts (continued)**

**Other information**  
■ store at 20-25°C (68-77°F)  
■ do not freeze  
■ do not use if printed neckband is broken or missing  
■ see bottom of box for lot number and expiration date

**Inactive ingredients**  
anhydrous citric acid, FD&C red #40, glycerin, hypromellose, natural and artificial berry flavor, polysorbate 80, purified water, sodium benzoate, sorbitol solution, sucrose, xanthan gum

**Questions or comments?**  
1-800-719-9260

**Gluten Free**

\*\*Major® Ibuprofen Oral Suspension 50 mg per 1.25 mL is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Motrin® Infants' Drops.

**Distributed by**  
**MAJOR®**  
**PHARMACEUTICALS**  
31778 Enterprise Drive  
Livonia, MI 48150 USA  
M-05 REV. 07/14  
Re-Order No. 242107



3 09045 46335 2

LOT NO.

EXP.

: 05705 5C C6

**CODE**  
**AREA**

