

Drug Facts

Active ingredient (in each 2 TSP (10 mL)) **Purpose**
Guaifenesin USP 200 mg.....Expectorant

Uses

helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive

Warnings

Ask a doctor before use if you have

- a persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- a cough occurs with too much phlegm (mucus)

Stop use and ask a doctor if cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of the reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than 6 doses in any 24 hour period
- this product is not intended for use in children under 12 years of age
- TSP = teaspoonful

adults and children 12 years and over	2 - 4 TSP every 4 hours
children under 12 years	do not use

Other information

- each TSP (5 mL) contains: **sodium 3 mg**
- store at 20°-25°C (68°-77°F). Do not refrigerate.

Inactive ingredients caramel powder, citric acid, corn syrup, FD&C Red #40, glycerin, menthol, propylene glycol, purified water, saccharin sodium, sodium benzoate

Questions or comments? (800) 616-2471

NDC 0904-0061-16
TAMPER-EVIDENT

RobafenTM Syrup

(brand of Guaifenesin syrup USP
100 mg per 5 mL)

EXPECTORANT

Each 2 teaspoonful (10 mL) contains:
Guaifenesin
(Glyceryl Guaiacolate), USP . . 200 mg

TAMPER-EVIDENT: Do not use this product if inner foil seal over the mouth of the bottle is cut, torn, broken or missing.

For Dispensing Purpose Only



Distributed by
MAJOR[®] PHARMACEUTICALS
17177 N Laurel Park Drive, Suite 233,
Livonia, MI 48152

NET: 1 Pint (473 mL)



M-64 Rev. 10/16 Re-order 140418,