


# SORBACT® COMPRESS

Infection preventive wound contact layer



## A WOUND CONTACT LAYER THAT REDUCES BIOBURDEN

*Sorbact® Compress is a sterile, bacteria and fungi binding wound dressing. It consists of a green Sorbact® wound contact layer, which allows passage of wound exudate into a secondary dressing. The dressing is folded into 8 layers. The dressing can be used in compression therapy.*

| Product           | Ref. no. | Size   | Size unfolded | Pcs/pack |
|-------------------|----------|--------|---------------|----------|
| Sorbact® Compress | 98128    | 4x6 cm | 11x15 cm      | 40       |
| Sorbact® Compress | 98125    | 7x9 cm | 17x28 cm      | 40       |



## INTENDED USE

Sorbact® Compress is a sterile, bacteria and fungi binding wound dressing. It consists of a green Sorbact® wound contact layer, which allows passage of wound exudate into a secondary dressing. The dressing is folded into 8 layers. The dressing can be used in compression therapy.

## INSTRUCTIONS FOR USE

1. Prepare the wound according to local clinical practice.
2. Select an appropriate dressing size for the wound. The dressing can be unfolded into suitable size and may overlap the wound margins if needed.
3. Remove the dressing from the pouch using an aseptic technique.
4. If cutting the dressing, use an aseptic technique. Discard any open and unused dressing.
5. Apply the dressing. Ensure that the dressing comes into direct contact with the complete wound surface. In cavity wounds and fistulas unfold and fluff up one or more dressings and line/fill the cavity or fistula. Avoid dense packing.
6. Apply an appropriate secondary dressing.
7. The dressing change frequency depends on exudate levels and the overall condition of the wound and surrounding skin. Should the clinical condition allow, the dressing can be left in place for up to 7 days.



## SORBACT® – INFECTION PREVENTIVE DRESSINGS

Sorbact® dressings employ a natural mechanism to bind microorganisms to their surface. The microorganisms are removed when the dressing is changed and the bioburden is thereby decreased.